

References

- Adgate JL, Barr DB, Clayton CA, Eberly LE, Freeman NC, Lioy PJ, et al. Measurement of children's exposure to pesticides: analysis of urinary metabolite levels in a probability-based sample. *Environ Health Perspect* 2001;109(6):583-90.
- Adlercreutz CH, Goldin BR, Gorbach SL, Hockerstedt KA, Watanabe S, Hamalainen EK, et al. Soybean phytoestrogens intake and cancer risk. *J Nutr* 1995a; 125(3 Suppl):757S-770s.
- Adlercreutz H, van der Wildt J, Kinzel J, Attalla H, Wahala K, Makela T, et al. Lignan and isoflavonoid conjugates in human urine. *J Steroid Biochem Mol Biol* 1995b;52(1) 97-103.
- Agency for Toxic Substances and Disease Registry. Toxicological Profiles (ATSDR) [online]. Available from URL: <http://www.atsdr.cdc.gov/toxpro2.html>, 9/30/02.
- Albertazzi P, Pansini F, Bottazi M, Bonaccorsi G, De Aloysio D, Morton MS. Dietary soy supplementation and phytoestrogen levels. *Obstet Gynecol* 1999;94(2):229-31.
- Alexandersson R. Blood and urinary concentrations as estimators of cobalt exposure. *Arch Environ Health* 1988;43(4):299-303.
- Allain P, Berre S, Premel-Cabic A, Mauras Y, Cledes A, Cournot A. Urinary elimination of molybdenum by healthy subjects as determined by inductively coupled plasma mass spectrometry. *Magnes Trace Elem* 1991-1992;10(1):47-50.
- Ambrose AM, Yost DH. Pharmacologic and toxicologic studies of N,N-diethyltoluamide. II. N,N-Diethyl-o-toluamide and N,N-diethyl-p-toluamide. *Toxicol Appl Pharmacol* 1965; 7(6):772-80.
- American Conference of Government Industrial Hygienists (ACGIH). 2000 TLVs and BEIs: threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati (OH): ACGIH, 2000.
- American Conference of Government Industrial Hygienists (ACGIH). Documentation of biological exposure indices. 7th edition. Cincinnati (OH); ACGIH Worldwide, 2001.
- Anderson HA, Falk C, Hanrahan L, Olson J, Burse VW, Needham LL, et al. Profiles of Great Lakes critical pollutants: a sentinel analysis of human blood and urine. The Great Lakes Consortium. *Environ Health Perspect* 1998;106(5):279-89.
- Anderson WA, Castle L, Scotter MJ, Massey RC, Springall C. A biomarker approach to measuring human dietary exposure to certain phthalate diesters. *Food Addit Contam* 2001;18(12):1068-74.
- Angerer J, Heinzel B, Reimann DO, Knorz W, Lehner G. Internal exposure to organic substances in a municipal waste incinerator. *Int Arch Occup Environ Health* 1992;64(4):265-73.
- Angerer J, Maass R, Heinrich R. Occupational exposure to hexachlorocyclohexane. VI. Metabolism of gamma-hexachlorocyclohexane in man. *Int Arch Occup Environ Health* 1983;52(1):59-67.
- Apostoli P, Schaller KH. Urinary beryllium—a suitable tool for assessing occupational and environmental beryllium exposure? *Int Arch Occup Environ Health* 2001;74(3):162-6.
- Aprea C, Sciarra G, Orsi D, Boccalon P, Sartorelli P, Sartorelli E. Urinary excretion of alkylphosphates in the general population (Italy). *Sci Total Environ* 1996;177(1-3):37-41.
- Aprea C, Strambi M, Novelli MT, Lunghini L, Bozzi N. Biologic monitoring of exposure to organophosphorus pesticides in 195 Italian children. *Environ Health Perspect* 2000;108(6):521-5.
- Arai Y, Uehara M, Sato Y, Kimura M, Eboshida A, Adlercreutz H. Comparison of isoflavones among dietary intake, plasma concentration and urinary excretion for accurate estimation of phytoestrogens intake. *J Epidemiol* 2000;10(2):127-35.
- Araki S, Sato H, Yokoyama K, Murata K. Subclinical neurophysiological effects of lead: A review on periph-

- eral, central, and autonomic nervous system effects in lead workers. *Am J Ind Med* 2000;37(2):193-204.
- Aylward LL, Hays SM. Temporal trends in human TCDD body burden: decreases over three decades and implications for exposure levels. *J Expo Anal Environ Epidemiol* 2002;12(5):319-28.
- Bailly R, Lauwerys R, Buchet JP, Mahieu P, Konings J. Experimental and human studies on antimony metabolism: their relevance for the biological monitoring of workers exposed to inorganic antimony. *Br J Ind Med* 1991;48(2):93-7.
- Baker DB, Loo S, Barker J. Evaluation of human exposure to the heptachlor epoxide contamination of milk in Hawaii. *Hawaii Med J* 1991;50(3):108-12, 118.
- Baker SE, Barr DB, Driskell WJ, Beeson MD, Needham LL. Quantification of selected pesticide metabolites in human urine using isotope dilution high-performance liquid chromatography/tandem mass spectrometry. *J Expo Anal Environ Epidemiol* 2000;10(6 Pt 2):789-98.
- Balluz L, Philen R, Ortega L, Rosales C, Brock J, Barr D, et al. Investigation of systemic lupus erythematosus in Nogales, Arizona. *Am J Epidemiol* 2001;154(11):1029-36.
- Bartels MJ, Kastl PE. Analysis of 3,5,6-trichloropyridinol in human urine using negative-ion chemical ionization gas chromatography-mass spectrometry. *J Chromatogr* 1992;575(1):69-74.
- Becker K, Kaus S, Krause C, Lepom P, Schulz C, Seiwert M, et al. German Environmental Survey 1998 (GerES III): Environmental pollutants in the blood of the German population. *Int J Hyg Environ Health* 2002;205(4):297-308.
- Benes B, Spevackova V, Smid J, Cejchanova M, Cerna M, Subrt P, et al. The concentration levels of Cd, Pb, Hg, Cu, Zn, and Se in blood of the population in the Czech Republic. *Cent Eur J Public Health* 2000;8(2):117-9.
- Bertram HP, Kemper FH, Muller C. Hexachlorobenzene content in human whole blood and adipose tissue: experiences in environmental specimen banking. *IARC Sci Publ* 1986; 77:173-82.
- Blount BC, Silva MJ, Caudill SP, Needham LL, Pirkle JL, Sampson EJ, et al. Levels of seven urinary phthalate metabolites in a human reference population. *Environ Health Perspect* 2000;108(10):979-82.
- Boffetta P, Hjourenkova N, Gustavsson P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes Control* 1997;8(3):444-72.
- Borja-Aburto VH, Hertz-Pannier I, Rojas LM, Farias P, Rios C, Blanco J. Blood lead levels measured prospectively and risk of spontaneous abortion. *Am J Epidemiol* 1999;150(6):590-7.
- Brockhaus A, Dolger R, Ewers U, Kramer U, Sodemann H, Wiegand H. Intake and health effects of thallium among a population living in the vicinity of a cement plant emitting thallium-containing dust. *Int Arch Occup Environ Health* 1981;48(4):375-89.
- Brokopp CD, Wyatt JI, Gabica J. Dialkyl phosphates in urine samples from pesticide formulators exposed to disulfoton and phorate. *Bull Environ Contam Toxicol* 1981;26(4):524-9.
- Budavari S, ed. *The Merck Index*. 11th ed. Rahway (NJ): Merck & Co., Inc;1989. p.308-9.
- Burns JE. Pesticides in people: organochlorine pesticide and polychlorinated biphenyl residues in biopsied human adipose tissue—Texas 1969-1972. *Pesticides Monitoring Journal* 1974;7:122-6.
- Byrne AR, Benedik L. Uranium content of blood, urine and hair of exposed and non-exposed persons determined by radiochemical neutron activation analysis, with emphasis on quality control. *Sci Total Environ* 1991;107:143-57.
- Byrne SL, Shurdut BA, Saunders DG. Potential chlorpyrifos exposure to residents following standard crack and crevice treatment. *Environ Health Perspect* 1998;106(11):725-31.
- Cabral J, Hall R, Rossi L, Bronczyk SA, Shubik P. Effects of long-term intake of DDT on rats. *Tumori* 1982; 68:11-7.

- Calvert GM, Wille KK, Sweeney MH, Fingerhut MA, Halperin WE. Evaluation of serum lipid concentrations among U.S. workers exposed to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Arch Environ Health* 1996;51(2):100-7.
- Caraballo R, Giovino G, Pechacek TF, Mowery PD, Richter PA, Strauss WJ, et al. Racial/ethnic differences in serum cotinine levels among adult U.S. cigarette smokers: The Third National Health and Nutrition Examination Survey, 1988-1991. *JAMA* 1998;280:135-40.
- Cassidy A, Bingham S, Setchell KD. Biological effects of a diet of soy protein rich in isoflavones on the menstrual cycle of premenopausal women. *Am J Clin Nutr* 1994;60(3):333-40.
- Center for the Evaluation of Risks to Human Reproduction of the National Toxicology Program. [online]. Available from URL: <http://cerhr.niehs.nih.gov/news>, 9/30/02.
- Centers for Disease Control and Prevention (CDC). Managing elevated blood lead levels among young children: recommendations from the Advisory Committee on Childhood Lead Poisoning Prevention. Atlanta (GA): CDC, 2002
- Christensen JM, Poulsen OM. A 1982-1992 surveillance programme on Danish pottery painters. Biological levels and health effects following exposure to soluble or insoluble cobalt compounds in cobalt blue dyes. *Sci Total Environ* 1994;50(1-3):95-104.
- Chuang JC, Callahan PJ, Lyu CW, Wilson NK. Polycyclic aromatic hydrocarbon exposures of children in low-income families. *J Expo Anal Environ Epidemiol* 1999;9(2):85-98.
- Clement J, Okey A. Reproduction in female rats born to DDT-treated parents. *Bull Environ Contam Toxicol* 1974;12(3):373-7.
- Dang HS, Pullat VR, Pillai KC. Determining the normal concentration of uranium in urine and application of the data to its biokinetics. *Health Phys* 1992;62(6):562-6.
- Das YT, Taskar PK, Brown HD, Chattopadhyay SK. Exposure of professional pest control operator to dichlorvos (DDVP) and residue on house structures. *Toxicol Lett* 1983;17(1-2):95-9.
- Davies JE, Peterson JC. Surveillance of occupational, accidental, and incidental exposure to organophosphate pesticides using urine alkyl phosphate and phenolic metabolite measurements. *Ann NY Acad Sci* 1997;837:257-68.
- Davies JE, Freed VH, Enos HF, Duncan RC, Barquet A, Morgade C, et al. Reduction of pesticide exposure with protective clothing for applicators and mixers. *J Occup Med* 1982; 24(6):464-8.
- den Tonkelaar I, Keinan-Boker L, Veer PV, Arts CJ, Adlercreutz H, Thijssen JH, et al. Urinary phytoestrogens and post menopausal breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2001;10(3):223-8.
- Deutsche Forschungsgemeinschaft (DFG). List of MAK and BAT values, 2000. Report No. 35, Commission for the Investigation of Health Hazards of Chemical Compounds in the work area, Weinheim, Wiley-VCH, 2000.
- Dewailly E, Ayotte P, Bruneau S, Lebel G, Levallois P, Weber JP. Exposure of the Inuit population of Nunavik (Arctic Quebec) to lead and mercury. *Arch Environ Health* 2001; 56(4):350-7.
- Dirven HA, van der Broek PH, Jongeneelen FJ. Determination of four metabolites of the plasticizer di (2-ethylhexyl) phthalate in human urine samples. *Int Arch Occup Environ Health* 1993;64(8):555-60.
- Dixon RA, Ferreira D. Genistein. *Phytochemistry* 2002;60(3):205-11.
- Dorgan JF, Brock JW, Rothman N, Needham LL, Miller R, Stephenson HE Jr, et al. Serum organochlorine pesticides and PCBs and breast cancer risk: results from a prospective analysis (USA). *Cancer Causes Control* 1999;10(1):1-11.
- Draper WM, Street JC. Applicator exposure to 2,4-D, dicamba, and a dicamba isomer. *J Environ Sci Health B* 1982;17(4):321-39.

- Dua VK, Pant CS, Sharma VP. Determination of levels of HCH and DDT in soil, water and whole blood from bioenvironmental and insecticide-sprayed areas of malaria control. Indian J Malariaol 1996;33(1):7-15.
- Duggan RE, Corneliusen PE. Dietary intake of pesticide chemicals in the United States. 3. June 1968–April 1970. Pestic Monit J 1972;5(4):331-41.
- Durham WF, Armstrong JF, Quinby GE. DDT and DDE content of complete prepared meals. Arch Environ Health 1965;11(5):641-7.
- Esteban E, Rubin C, Hill R, Olson D, Pearce K. Association between indoor residential contamination with methyl parathion and urinary *para*-nitrophenol. J Expo Anal Environ Epidemiol 1996;6(3):375-87.
- Ejnik JW, Carmichael AJ, Hamilton MM, McDiarmid M, Squibb K, Boyd P, et al. Determination of the isotopic composition of uranium in urine by inductively coupled plasma mass spectrometry. Health Phys 2000; 78(2):143-6.
- Eriksson P, Archer T, Fredriksson A. Altered behaviour in adult mice exposed to a single low dose of DDT and its fatty acid conjugate as neonates. Brain Res 1990;514(1):141-2.
- Ewers U, Krause C, Schulz C, Wilhelm M. Reference values and human biological monitoring values for environmental toxins. Report on the work and recommendations of the Commission on Human Biological Monitoring of the German Federal Environmental Agency. Int Arch Occup Environ Health 1999; 72(4):255-60.
- Fabro S, McLachlan JA, Dames NM. Chemical exposure of embryos during the preimplantation stages of pregnancy: mortality rate and intrauterine development. Am J Obstet Gynecol 1984;148(7):929-38.
- Falk C, Hanrahan L, Anderson HA, Kanarek MS, Draheim L, Needham LL. Body burden levels of dioxin, furans, and PCBs among frequent consumers of Great Lakes sport fish. The Great Lakes Consortium. Environ Res 1999;80(2 Pt 2):S19-S25.
- Farago ME, Kavanagh P, Blanks R, Kelly J, Kazantzis G, Thornton I, et al. Platinum concentrations in urban road dust and soil, and in blood and urine in the United Kingdom. Analyst 1998;123(3):451-4.
- Fenske RA, Lu C, Barr D, Needham L. Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. Environ Health Perspect 2002;110(5):549-53.
- Finklea J, Priester LE, Creason JP, Hauser T, Hinners T, Hammer DI. Polychlorinated biphenyl residues in human plasma expose a major urban pollution problem. Am J Public Health 1972; 62(5):645-51.
- Franklin CA, Fenske RA, Greenhalgh R, Mathieu L, Denley HV, Leffingwell JT, et al. Correlation of urinary pesticide metabolite excretion with estimated dermal contact in the course of occupational exposure to Guthion. J Toxicol Environ Health 1981;7(5):715-31.
- Geltman PL, Brown MJ, Cochran J. Lead poisoning among refugee children resettled in Massachusetts, 1995 to 1999. Pediatrics 2001;108(1):158-62.
- Geyer HJ, Schramm KW, Feicht EA, Behechi A, Steinberg C, Bruggemann R, et al. Half-lives of tetra-, penta-, hexa-, hepta-, and octachlorodibenzo-*p*-dioxin in rats, monkeys, and humans—a critical review. Chemosphere 2002;48(6):631-44.
- Glynn AW, Wolk A, Aune M, Atuma S, Zettermark S, Maehle-Schmid M, et al. Serum concentrations of organochlorines in men: a search for markers of exposure. Sci Total Environ 2000; 263(1-3),197-208.
- Goen T, Gundel J, Schaller KH, Angerer J. The elimination of 1-hydroxypyrene in the urine of the general population and workers with different occupational exposures to PAH. Sci Total Environ 1995;163(1-3):195-201.
- Grandjean P, Weihe P, Needham LL, Burse VW, Patterson DG Jr, Sampson EJ, et al. Relation of a seafood diet to mercury, selenium, arsenic, and polychlorinated biphenyl and other organochlorine concentrations in human milk. Environ Res 1995;71(1):29-38.

- Grimalt JO, Sunyer J, Moreno V, Amaral OC, Sala M, Rosell A, et al. Risk excess of soft-tissue sarcoma and thyroid cancer in a community exposed to airborne organochlorinated compound mixtures with a high hexachlorobenzene content. *Int J Cancer* 1994; 56(2):200-3.
- Gundel J, Mannschreck C, Buttner K, Ewers U, Angerer J. Urinary levels of 1-hydroxypyrene, 1-, 2-, 3-, and 4-hydroxyphenanthrene in females living in an industrial area of Germany. *Arch Environ Contam Toxicol* 1996;31(4):585-90.
- Gundel J, Schaller KH, Angerer J. Occupational exposure to polycyclic aromatic hydrocarbons in a fireproof stone producing plant: biological monitoring of 1-hydroxypyrene, 1-, 2-, 3- and 4-hydroxyphenanthrene, 3-hydroxybenz(a)anthracene and 3-hydroxybenzo-(a)-pyrene. *Int Arch Occup Environ Health* 2000;73(4):270-4.
- Halperin W, Vogt R, Sweeney MH, Shopp G, Fingerhut M, Petersen M. Immunological markers among workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Occup Environ Med* 1998;55(11):742-9.
- Hamilton EI, Sabbioni E, Van der Venne MT. Element reference values in tissues from inhabitants of the European Community. VI. Review of elements in blood, plasma and urine and a critical evaluation of reference values for the United Kingdom population. *Sci Total Environ* 1994;158(1-3):165-90.
- Hanrahan LP, Falk C, Anderson HA, Draheim L, Kanarek MS, Olson J. Serum PCB and DDE levels of frequent Great Lakes sport fish consumers—a first look. The Great Lakes Consortium. *Environ Res* 1999;80(2 Pt 2):S26-S37.
- Hayes W J, Jr. Mortality in 1969 from pesticides, including aerosols. *Arch Environ Health* 1976;31:61-72.
- Helzlsouer KJ, Alberg AJ, Huang HY, Hoffman SC, Strickland PT, Brock JW, et al. Serum concentrations of organochlorine compounds and the subsequent development of breast cancer. *Cancer Epidemiol Biomarkers Prev* 1999;8(6):525-32.
- Hernandez-Valero MA, Bondy ML, Spitz MR, Zahm SH. Evaluation of Mexican American migrant farm-worker work practices and organochlorine pesticide metabolites. *Am J Ind Med* 2001; 40(5):554-60.
- Herrero C, Ozalla D, Sala M, Otero R, Santiago-Silva M, Lecha M, et al. Urinary porphyrin excretion in a human population highly exposed to hexachlorobenzene. *Arch Dermatol* 1999;135(4):400-4.
- Heudorf U, Angerer J. Internal exposure to PAHs of children and adults living in homes with parquet flooring containing high levels of PAHs in the parquet glue. *Int Arch Occup Environ Health* 2001;74(2):91-101.
- Hill RH Jr, Head SL, Baker S, Gregg M, Shealy DB, Bailey SL, et al. Pesticide residues in urine of adults living in the United States: reference range concentrations. *Environ Res* 1995;71(2):99-108.
- Hill RH Jr, To T, Holler JS, Fast DM, Smith SJ, Needham LL, et al. Residues of chlorinated phenols and phenoxy acid herbicides in the urine of Arkansas children. *Arch Environ Contam Toxicol* 1989;18(4):469-74.
- Hodgert JH, Zacharewski TR, Hammond GL. Interactions between human plasma sex hormone-binding globulin and xenobiotic ligands. *J Steroid Biochem Mol Biol* 2000;75(2-3):167-76.
- Hoppin JA, Brock JW, Davis BJ, Baird DD. Reproducibility of urinary phthalate metabolites in first morning urine samples. *Environ Health Perspect* 2000;110(5):515-8.
- Horn-Ross PL, Barnes S, Kirk M, Coward L, Parsonnet J, Hiatt RA. Urinary phytoestrogen levels in young women from a multiethnic population. *Cancer Epidemiol Biomarkers Prev* 1997;6(5):339-45.
- Hoyer AP, Grandjean P, Jorgensen T, Brock JW, Hartvig HB. Organochlorine exposure and risk of breast cancer. *Lancet* 1998;352(9143):1816-20.
- Hunter DJ, Hankinson SE, Laden F, Colditz GA, Manson JE, Willett WC, et al. Plasma organochlorine levels and the risk of breast cancer. *N Engl J Med* 1997;337(18):1253-8.

- Hutchins AM, Slavin JL, Lampe JW. Urinary isoflavanoid phytoestrogen and lignan excretion after consumption of fermented and unfermented soy products. *J Am Diet Assoc* 1995;95(5):545-51.
- Ichikawa Y, Kusaka Y, Goto S. Biological monitoring of cobalt exposure, based on cobalt concentrations in blood and urine. *Int Arch Occup Environ Health* 1985;55(4):269-76.
- Ingram D, Sanders K, Kolybaba M, Lopez D. Case-control study of phyto-oestrogens and breast cancer. *Lancet* 1997;350(9083):990-4.
- Institute of Medicine. Veterans and Agent Orange-Update 2000. Washington D.C., National Academy Press, 2001. p.5-12.
- International Agency for Research on Cancer (IARC), World Health Organization. IARC Monographs. [online]. Available from URL: <http://www.iarc.fr/>, 9/30/02.
- International Programme on Chemical Safety. Environmental Health Criteria Monographs. Carbamate pesticides: a general introduction. Vol 64, World Health Organization. 1986, p.23-43.
- Iversen BS, Menne C, White MA, Kristiansen J, Christensen JM, Sabbioni E. Inductively coupled plasma mass spectrometric determination of molybdenum in urine from a Danish population. *Analyst* 1998;123(1):81-5.
- Jauhainen A, Kangas J, Laitinen S, Savolainen K. Biological monitoring of workers exposed to mevinphos in greenhouses. *Bull Environ Contam Toxicol* 1992; 49(1):37-43.
- Jemal A, Graubard BI, Devesa SS, Flegel KM. The association of blood lead level and cancer mortality among whites in the United States. *Environ Health Perspect* 2002;110(4):325-9.
- Johnson GD, Harsy SG, Geronimo J, Wise JM. Orthophenylphenol and phenylhydroquinone residues in citrus fruit and processed citrus products after postharvest fungicidal treatments with sodium orthophenylphenate in California and Florida. *J Agric Food Chem* 2001;49(5):2497-502.
- Jones RD, Winter DP, Cooper AJ. Absorption study of pentachlorophenol in persons working with wood preservatives. *Hum Toxicol* 1986;5(3):189-94.
- Jongeneelen FJ. Biological monitoring of environmental exposure to polycyclic aromatic hydrocarbons; 1-hydroxypyrene in urine of people. *Toxicol Lett* 1994;72(1-3):205-11.
- Jonsson HT Jr, Keil JE, Gaddy RG, Loadholt CB, Hennigar GR, Walker EM Jr. Prolonged ingestion of commercial DDT and PCB; effects on progesterone levels and reproduction in the mature female rat. *Arch Environ Contam Toxicol* 1975-76;3(4):479-90.
- Jung D, Berg PA, Edler L, Ehrenthal W, Fenner D, Flesch-Janys D, et al. Immunologic findings in workers formerly exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin and its congeners. *Environ Health Perspect* 1998;106(Suppl 2):689-95.
- Kaiser R, Henderson AK, Daley WR, Naughton M, Khan MH, Rahman M, et al. Blood lead levels of primary school children in Dhaka, Bangladesh. *Environ Health Perspect* 2001;109(6):563-6.
- Kang D, Tepper A, Patterson DG Jr. Coplanar PCBs and the relative contribution of coplanar PCBs, PCDDs, and PCDFs to the total 2,3,7,8-TCDD toxicity equivalents in human serum. *Chemosphere* 1997;35(3):503-11.
- Kanoh T, Fukuda M, Onozuka H, Kinouchi T, Ohnishi Y. Urinary 1-hydroxypyrene as a marker of exposure to polycyclic aromatic hydrocarbons in environment. *Environ Res* 1993;62(2):230-41.
- Karpas Z, Halicz L, Roiz J, Marko R, Katorza E, Lorber A, et al. Inductively coupled plasma mass spectrometry as a simple, rapid, and inexpensive method for determination of uranium in urine and fresh water: comparison with LIF. *Health Phys* 1996;71(6):879-85.
- Karpas Z, Lorber A, Elish E, Kol R, Roiz Y, Marko R, et al. Uptake of ingested uranium after low "acute intake." *Health Phys* 1998;74(3):337-45.
- Karr SC, Lampe JW, Hutchins AM, Slavin JL. Urinary isoflavanoid excretion in humans is dose dependent at low to moderate levels of soy-protein consumption. *Am J Clin Nutr* 1997;66(1):46-51.

- Kentner M, Leinemann M, Schaller KH, Weltle D, Lehnert G. External and internal antimony exposure in starter battery production. *Int Arch Occup Environ Health* 1995;67(2):119-23.
- Kim MK, Chung BC, Yu VY, Nam JH, Lee HC, Huh KB, et al. Relationships of urinary phyto-oestrogen excretion to BMD in postmenopausal women. *Clin Endocrinol (Oxf)* 2002;156(3):321-8.
- Knowles LM, Zigrossi DA, Tauber RA, Hightower C, Milner JA. Flavonoids suppress androgen-independent human prostate tumor proliferation. *Nutr Cancer* 2000;38(1):116-22.
- Kolmodin-Hedman B, Erne K. Estimation of occupational exposure to phenoxy acids (2,4-D and 2,4,5-T). *Arch Toxicol Suppl* 1980;4:318-21.
- Komaromy-Hiller G, Ash KO, Costa R, Howerton K. Comparison of representative ranges based on U.S. patient population and literature reference intervals for urinary trace elements. *Clin Chim Acta* 2000;296(1-2):71-90.
- Koo JW, Parham F, Kohn MC, Masten SA, Brock JW, Needham LL, et al. The association between biomarker-based exposure estimates for phthalates and demographic factors in a human reference population. *Environ Health Perspect* 2002;110(4):405-10.
- Koopman-Esseboom C, Huisman M, Touwen BC, Boersma ER, Brouwer A, Sauer PJ, et al. Newborn infants diagnosed as neurologically abnormal with relation to PCB and dioxin exposure and their thyroid-hormone status. *Dev Med Child Neurol* 1997;39(11):785.
- Korn EL, Graubard BI. Confidence intervals for proportions with small expected number of positive counts estimated from survey data. *Survey Methodology* 1998; 24:193-201.
- Kotsopoulos D, Dalais FS, Liang YL, McGrath BP, Teede HJ. The effects of soy protein-containing phytoestrogens on menopausal symptoms in postmenopausal women. *Climacteric* 2000;3(3):161-7.
- Kraus T, Schramel P, Schaller KH, Zobelein P, Weber A, Angerer J. Exposure assessment in the hard metal manufacturing industry with special regard to tungsten and its compounds. *Occup Environ Med* 2001;58(10):631-4.
- Kurttio P, Auvinen A, Salonen L, Saha H, Pekkanen J, Makelainen I, et al. Renal effects of uranium in drinking water. *Environ Health Perspect* 2002;110(4):337-42.
- Kutz FW, Yobs AR, Strassman SC. Racial stratification of organochlorine insecticide residues in human adipose tissue. *J Occup Med* 1977;19(9):619-22.
- Kutz FW, Wood PH, Bottimore DP. Organochlorine pesticides and polychlorinated biphenyls in human adipose tissue. *Rev Environ Contam Toxicol* 1991;120:1-82.
- Lampe JW, Gustafson DR, Hutchins AM, Martini MC, Li S, Wahala K, et al. Urinary isoflavonoid and lignan excretion on a Western diet: relation to soy, vegetable, and fruit intake. *Cancer Epidemiol Biomarkers Prev* 1999;8(8):699-707.
- Lauwerys RB, Hoet P. Industrial Chemical Exposure: Guidelines for Biological Monitoring. 3rd ed. Boca Raton (FL): Lewis Publishers, 2001. p. 87-95, 54-68, and 500-11.
- Lebel G, Dodin S, Ayotte P, Marcoux S, Ferron LA, Dewailly E. Organochlorine exposure and the risk of endometriosis. *Fertil Steril* 1998;69(2):221-8.
- Leng G, Lewalter J. Role of individual susceptibility in risk assessment of pesticides. *Occup Environ Med* 1999;56(7):449-53.
- Linnainmaa M, Kiilunen M. Urinary cobalt as a measure of exposure in the wet sharpening of hard metal and stellite blades. *Int Arch Occup Environ Health* 1997;69(3):193-200.
- Lison D, Buchet JP, Swennen B, Molders J, Lauwerys R. Biological monitoring of workers exposed to cobalt metal, salt, oxides, and hard metal dust. *Occup Environ Med* 1994;51(7):447-50.
- Liss GM, Albro PW, Hartle RW, Stringer WT. Urine phthalate determinations as an index of occupational exposure to phthalic anhydride and di (2-ethylhexyl) phthalate. *Scand J Work Environ Health* 1985;11(5):381-7.

- Lloyd J. Long-term mortality study of steelworkers. V. Respiratory cancer in coke plant workers. *J Occup Med* 1971;13:53-68.
- Longnecker MP, Ryan JJ, Gladen BC, Schecter AJ. Correlations among human plasma levels of dioxin-like compounds and polychlorinated biphenyls (PCBs) and implications for epidemiologic studies. *Arch Environ Health* 2000;55(3):195-200.
- Lu LJ, Grady JJ, Marshall MV, Ramanujam VM, Anderson KE. Altered time course of urinary daidzein and genistein excretion during chronic soya diet in healthy male subjects. *Nutr Cancer* 1995;24(3):311-23.
- Ludersdorf R, Fuchs A, Mayer P, Skulsuksai G, Schacke G. Biological assessment of exposure to antimony and lead in the glass-producing industry. *Int Arch Occup Environ Health* 1987;59(5):469-74.
- Lundberg C. Effect of DDT on cytochrome P-450 and oestrogen-dependent functions in mice. *Environ Physiol Biochem* 1974; 4:200-4.
- Luotamo M, Jarvisalo J, Aitio A. Assessment of exposure to polychlorinated biphenyls: analysis of selected isomers in blood and adipose tissue. *Environ Res* 1991;54(2):121-34.
- MacIntosh DL, Needham LL, Hammerstrom KA, Ryan PB. A longitudinal investigation of selected pesticide metabolites in urine. *J Expo Anal Environ Epidemiol* 1999;9(5):494-501.
- Mahaffey KR, Mergler D. Blood levels of total and organic mercury in residents of the upper St. Lawrence River basin, Quebec: association with age, gender, and fish consumption. *Environ Res* 1998;77(2):104-14.
- McRill C, Boyer LV, Flood TJ, Ortega L. Mercury toxicity due to use of a cosmetic cream. *J Occup Environ Med* 2000;42(1):4-7.
- Marcus RL. Investigation of a working population exposed to thallium. *J Soc Occup Med* 1985;35(1):4-9.
- Masuda Y. Fate of PCDF/PCB congeners and change of clinical symptoms in patients with Yusho PCB poisoning for 30 years. *Chemosphere* 2001 43(4-7):925-30.
- Masuda Y, Schecter A, Papke O. Concentrations of PCBs, PCDFs and PCDDs in the blood of Yusho patients and their toxic equivalent contribution. *Chemosphere* 1998;37(9-12):1773-80.
- Matsuura N, Uchiyama T, Tada H, Nakamura Y, Kondo N, Morita M. Effects of dioxins and polychlorinated biphenyls (PCBs) on thyroid function in infants born in Japan—the second report from research on environmental health. *Chemosphere* 2001;45(8):1167-71.
- Melnick RL. Is peroxisome proliferation an obligatory precursor step in the carcinogenicity of di (2-ethylhexyl) phthalate (DEHP)? *Environ Health Perspect* 2001;109(5):437-42.
- Michalek JE, Pirkle JL, Needham LL, Patterson DG Jr, Caudill SP, Tripathi RC, et al. Pharmacokinetics of 2,3,7,8-tetrachlorodibenzo-p-dioxin in Seveso adults and veterans of Operation Ranch Hand. *J Expo Anal Environ Epidemiol* 2002;12(1):44-53.
- Michalek JE, Akhtar FZ, Kiel JL. Serum dioxin, insulin, fasting glucose, and sex hormone-binding globulin in veterans of Operation Ranch Hand. *J Clin Endocrinol Metab* 1999;84(5):1540-3.
- Minoia C, Sabbioni E, Apostoli P, Pietra R, Pozzoli L, Gallorini M, et al. Trace element reference values in tissues from inhabitants of the European community. I. A study of 46 elements in urine, blood and serum of Italian subjects. *Sci Total Environ* 1990;95:89-105.
- Mocarelli P, Needham LL, Marocchi A, Patterson DG Jr, Brambilla P, Gerthoux PM, et al. Serum concentrations of 2,3,7,8-tetrachlorodibenzo-p-dioxin and test results from selected residents of Seveso, Italy. *J Toxicol Environ Health* 1991;32(4):357-66.
- Morgan DP, Hetzler HL, Slach EF, Lin LI. Urinary excretion of paranitrophenol and alkyl phosphates following ingestion of methyl or ethyl parathion by human subjects. *Arch Environ Contam Toxicol* 1977;6(2-3):159-73.
- Murkies A, Dalais FS, Briganti EM, Burger HG, Healy DL, Wahlgqvist ML. Phytoestrogens and breast cancer in postmenopausal women: a case control study. *Menopause* 2000;7(5):289-96.

National Research Council. Toxicologic effects of methylmercury. Washington D.C., National Academy of Sciences, 2000.

National Research Council (Platinum Group Metals). Medical and biologic effects of environmental pollutants. Washington D.C., National Academy of Sciences, 1977. p. 232.

National Toxicology Program. Ninth Report on Carcinogens (NTP 9th ROC). U.S. Department of Health and Human Services, Public Health Service [online]. Available from URL: <http://ehis.niehs.nih.gov/roc/toc9.html>, 9/30/02.

National Toxicology Program. TR-301, Toxicology and carcinogenesis studies of *o*-phenylphenol alone and with 7,12-dimethylbenz(a)anthracene in Swiss CD-1 mice (dermal studies). TR-301 (NTIS # PB86-217239), March 1986.

Needham LL, Burse VW, Head SL, Korver MP, McClure PC, Andrews JS Jr, et al. Adipose tissue/serum partitioning of chlorinated hydrocarbon pesticides in humans. Chemosphere 1990; 20:975-80.

Nicholls J, Lasley BL, Nakajima ST, Setchell KD, Schneeman BO. Effects of soy consumption on gonadotropin secretion and acute pituitary responses to gonadotropin-releasing hormone in women. J Nutr 2002;132(4):708-14.

Nicolaou G, Pietra R, Sabbioni E, Mosconi G, Cassina G, Seghizzi P. Multielement determination of metals in biological specimens of hard-metal workers: a study carried out by neutron activation analysis. J Trace Elem Electrolytes Health Dis 1987;1(2):73-7.

Nigam SK, Karnik AB, Majumder SK, Visweswariah K, Raju GS, Bai KM, et al. Serum hexachlorocyclohexane residues in workers engaged at a HCH manufacturing plant. Int Arch Occup Environ Health. 1986;57(4):315-20.

Occupational Safety and Health Administration. 29 CFR 1910.1127 Section 1-Medical Surveillance. Washington (D.C.): OSHA, 1992.

Occupational Safety and Health Administration (OSHA) [online]. Available from URL: <http://www.osha.gov/comp-links.html>, 9/30/02.

Papke O. PCDD/PCDD: human background data for Germany, a 10-year experience. Environ Health Perspect 1998;106 (Suppl 2):723-31.

Parkinson A, Safe SH, Robertson LW, Thomas PE, Ryan DE, Reik LM, et al. Immunochemical quantitation of cytochrome P-450 isozymes and epoxide hydrolase in liver microsomes from polychlorinated or polybrominated biphenyl-treated rats. A study of structure-activity relationships. J Biol Chem 1983;258(9):5967-76.

Paschal DC, Burse V, Gunter EW, Pirkle JL, Sampson EJ, Miller DT, et al. Exposure of the U.S. population aged 6 years and older to cadmium: 1988-1994. Arch Environ Contam Toxicol 2000;38:377-83.

Paschal DC, Ting BG, Morrow JC, Pirkle JL, Jackson RJ, Sampson EJ, et al. Trace metals in urine of United States residents: reference range concentrations. Environ Res 1998;76(1):53-9.

Patterson DG Jr, Todd GD, Turner WE, Maggio V, Alexander LR, Needham LL. Levels of non-ortho-substituted (coplanar), mono- and di-ortho-substituted polychlorinated biphenyls, dibenzo-*p*-dioxins, and dibenzofurans in human serum and adipose tissue. Environ Health Perspect 1994;102 (Suppl 1):195-204.

Peck CC, Albro PW. Toxic potential of the plasticizer di (2-ethylhexyl) phthalate in the context of its disposition and metabolism in primates and man. Environ Health Perspect 1982;45:11-7.

Pesch A, Wilhelm M, Rostek U, Schmitz N, Weishoff-Houben M, Ranft U, et al. Mercury concentrations in urine, scalp hair, and saliva in children from Germany. J Expo Anal Environ Epidemiol 2002;12(4):252-8.

Peters HA, Gocmen A, Cripps DJ, Bryan GT, Dogramaci I. Epidemiology of hexachlorobenzene-induced porphyria in Turkey: clinical and laboratory follow-up after 25 years. Arch Neurol 1982;39(12):744-9.

Pirkle JL, Flegal KM, Bernert JT, Brody DJ, Etzel RA, Maurer KR. Exposure of the U.S. population to environmental tobacco smoke: the Third National Health and Nutrition Examination Survey, 1988-1991. JAMA 1996;275:1233-40.

- Pirkle JL, Kaufmann RB, Brody DJ, Hickman T, Gunter EW, Paschal DC. Exposure of the U.S. population to lead: 1991-1994. *Environ Health Perspect* 1998;106:745-50.
- Radomski JL, Astolfi E, Deichmann WB, Rey AA. Blood levels of organochlorine pesticides in Argentina: occupationally and nonoccupationally exposed adults, children and newborn infants. *Toxicol Appl Pharmacol* 1971;20(2):186-93.
- Redmond CK, Strobino BR, Cypess RH. Cancer experience among coke by-product workers. *Ann N Y Acad Sci* 1976;271:102-15.
- Richter ED, Kowalski M, Leventhal A, Grauer F, Marzouk J, Brenner S, et al. Illness and excretion of organophosphate metabolites four months after household pest extermination. *Arch Environ Health* 1992; 47(2):135-8.
- Ritchie KA, Gilmour WH, Macdonald EB, Burke FJ, McGowan DA, Dale IM, et al. Health and neuropsychological functioning of dentists exposed to mercury. *Occup Environ Med* 2002;59(5):287-93.
- Robbins PJ, Cherniack MG. Review of the biodistribution and toxicity of the insect repellent N,N-diethyl-m-toluamide (DEET). *J Toxicol Environ Health* 1986;18(4):503-25.
- Roels HA, Hoet P, Lison D. Usefulness of biomarkers of exposure to inorganic mercury, lead, or cadmium in controlling occupational and environmental risks of nephrotoxicity. *Ren Fail* 1999; 21(3-4):251-62
- Roggi C, Minoia C, Sciarra GF, Apostoli P, Maccarini L, Magnaghi S, et al. Urinary 1-hydroxypyrene as a marker of exposure to pyrene: an epidemiological survey on a general population group. *Sci Total Environ* 1997;199(3):247-54.
- Safe SH, Pallaroni L, Yoon K, Gaido K, Ross S, Saville B, et al. Toxicology of environmental estrogens. *Reprod Fertil Dev* 2001;13(4):307-15.
- Sala M, Sunyer J, Otero R, Santiago-Silva M, Camps C, Grimalt J. Organochlorine in the serum of inhabitants living near an electrochemical factory. *Occup Environ Med* 1999;56(3):152-8.
- Sanzo JM, Dorronsoro M, Amiano P, Amurrio A, Aguinagalde FX, Azpíri MA. Estimation and validation of mercury intake associated with fish consumption in an EPIC cohort of Spain. *Public Health Nutr* 2001; 4(5):981-8.
- SAS Institute Inc., SAS/STAT User's Guide, Version 8, Cary, NC: SAS Institute Inc.; 1999.
- Satarug S, Baker JR, Reilly PE, Moore MR, Williams DJ. Cadmium levels in the lung, liver, kidney cortex, and urine samples from Australians without occupational exposure to metals. *Arch Environ Health* 2002;57(1):69-77.
- Schaller KH, Manke G, Raithel HJ, Buhlmeyer G, Schmidt M, Valentin H. Investigations of thallium-exposed workers in cement factories. *Int Arch Occup Environ Health* 1980;47(3):223-31.
- Schierl R. Urinary platinum levels associated with dental gold alloys. *Arch Environ Health* 2001;56(3):283-6.
- Schierl R, Fries HG, van de Weyer C, Fruhmann G. Urinary excretion of platinum from platinum-industry workers. *Occup Environ Med* 1998;55(2):138-4.
- Schramel P, Wendler I, Angerer J. The determination of metals (antimony, bismuth, lead, cadmium, mercury, palladium, platinum, tellurium, thallium, tin and tungsten) in urine samples by inductively coupled plasma-mass spectrometry. *Int Arch Occup Environ Health* 1997;69(3):219-23.
- Seifert B, Becker K, Helm D, Krause C, Schulz C, Seiwert M. The German Environmental Survey 1990/1992 (GerES II): reference concentrations of selected environmental pollutants in blood, urine, hair, house dust, drinking water and indoor air. *J Expo Anal Environ Epidemiol* 2000; 10(6 Pt 1):552-65.
- Setchell KD, Brown NM, Desai P, Zimmer-Nechemias L, Wolfe BE, Brashear WT, et al. Bioavailability of pure isoflavones in healthy humans and analysis of commercial soy isoflavone supplements. *J Nutr* 2001;131(4 Suppl):1362S-75S.

- Sievers E, Schleyerbach U, Schaub J. Molybdenum in infancy: methodical investigation of urinary excretion. *J Trace Elem Med Biol* 2001;15(2-3):149-54.
- Simpson GR, Higgins V, Chapman J, Bermingham S. Exposure of council and forestry workers to 2,4,5-T. *Med J Aust* 1978;18;2(11):536-7.
- Slavin JL, Karr SC, Hutchins AM, Lampe JW. Influence of soybean processing, habitual diet, and soy dose on urinary isoflavonoid excretion. *Am J Clin Nutr* 1998;68(6 Suppl):1492S-1495S.
- Smallwood AW, DeBord KE, Lowry LK. N,N'-diethyl-m-toluamide (m-DET): analysis of an insect repellent in human urine and serum by high-performance liquid chromatography. *J Anal Toxicol* 1992;16(1):10-3.
- Stattin P, Adlercreutz H, Tenkanen L, Jellum E, Lumme S, Hallmans G, et al. Circulating enterolactone and prostate cancer risk: a Nordic nested case-control study. *Int J Cancer* 2002;99(1):124-9.
- Stehr-Green PA. Demographic and seasonal influences on human serum pesticide residue levels. *J Toxicol Environ Health* 1989;27(4):405-21.
- Sturgeon SR, Brock JW, Potischman N, Needham LL, Rothman N, Brinton LA, et al. Serum concentrations of organochlorine compounds and endometrial cancer risk (United States). *Cancer Causes Control* 1998;9(4):417-24.
- SUDAAN User's Manual, Release 8.0, Research Triangle Park, NC: Research Triangle Institute; 2001.
- Swennen B, Buchet JP, Stanescu D, Lison D, Lauwerys R. Epidemiological survey of workers exposed to cobalt oxides, cobalt salts, and cobalt metal. *Br J Ind Med* 1993;50(9):835-42.
- Szániszlo J, Ungvary G. Polycyclic aromatic hydrocarbon exposure and burden of outdoor workers in Budapest. *J Toxicol Environ Health A* 2001;62(5):297-306.
- Takamiya K. Monitoring of urinary alkyl phosphates in pest-control operators exposed to various organophosphorus insecticides. *Bull Environ Contam Toxicol* 1994;52(2):190-5.
- Taylor JK. Quality Assurance of Chemical Measurements. Chelsea, MI: Lewis Publishing, 1987.
- Teede HJ, Dalais FS, Kotsopoulos D, Liang YL, Davis S, McGrath BP. Dietary soy has both beneficial and potentially adverse cardiovascular effects: a placebo-controlled study in men and postmenopausal women. *J Clin Endocrinol Metab* 2001;86(7):3053-60.
- Tepper A, Burt S, Piacitelli L, Patterson DG Jr. Coplanar PCBs and the relative contribution of coplanar PCBs, PCDDs, and PCDFs to the total 2,3,7,8-TCDD toxicity equivalents in human serum. *Chemosphere* 1997;35(3):503-11.
- Ting BG, Paschal DC, Jarrett JM, Pirkle JL, Jackson RJ, Sampson EJ, et al. Uranium and thorium in urine of United States residents: reference range concentrations. *Environ Res* 1999;81(1):45-51.
- Tsongas TA, Meglen RR, Walravens PA, Chappell WR. Molybdenum in the diet: an estimate of average daily intake in the United States. *Am J Clin Nutr* 1980;33(5):1103-7.
- Turnlund JR, Keyes WR, Peiffer GL. Molybdenum absorption, excretion, and retention studied with stable isotopes in young men at five intakes of dietary molybdenum. *Am J Clin Nutr* 1995 Oct;62(4):790-6.
- U.S. Environmental Protection Agency (U.S. EPAa), Office of Pesticide Programs. Organophosphate pesticides in food: a primer on reassessment of residue limits [online]. Available from URL: <http://www.epa.gov/pesticides/op/primer.htm>, 9/30/02.
- U.S. Environmental Protection Agency (U.S. EPAb), Office of Pesticide Programs. Status summary of the organophosphate review process [online]. Available from URL: <http://www.epa.gov/pesticides/op/status.htm>, 9/30/02.
- U.S. Environmental Protection Agency (U.S. EPAc). [online]. Available from URL: <http://www.epa.gov/pesticides/citizens/deet.htm>, 9/30/02.
- U.S. Environmental Protection Agency (U.S. EPAd). Office of Prevention, Pesticides Reregistration Eligibility Decision (RED) DEET, EPA738-R-98-010, EPA: Washington, DC. September 1998.

- U.S. Environmental Protection Agency (EPA IRIS). Integrated Risk Information System [online]. Available from URL: <http://www.epa.gov/subst/index.html>, 9/30/02.
- U.S. Nuclear Regulatory Commission. U.S. Nuclear Regulatory Commission (NRC) Guide 8.22-Bioassay at Uranium Mills. Washington (D.C.): NRC; July 1978.
- van Wijnen JH, Slob R, Jongmans-Liedekerken G, van de Weerd RH, Woudenberg F. Exposure to polycyclic aromatic hydrocarbons among Dutch children. *Environ Health Perspect* 1996;104(5):530-4.
- Vaughan GT, Florence TM. Platinum in the human diet, blood, hair and excreta. *Sci Total Environ* 1992;111(1):47-58.
- Wariishi M, Suzuki Y, Nishiyama K. Chlordane residues in normal human blood. *Bull Environ Contam Toxicol* 1986; 36(5):635-43.
- Waluszewski SM, Pardio VT, Chantiri JN, Infanzon RM, Rivera J. Organochlorine pesticide residues in adipose tissue of Mexicans. *Sci Total Environ* 1996;181(2):125-31.
- Walker KC, Goette MB, Batchelor GS. Pesticide residues in foods: dichlorodiphenyltrichloroethane and dichlorodiphenyldichloroethylene content of prepared meals. *J Agric Food Chem* 1954;2:1034-7.
- White MA, Sabbioni E. Trace element reference values in tissues from inhabitants of the European Union. X. A study of 13 elements in blood and urine of a United Kingdom population. *Sci Total Environ* 1998; 216(3):253-70.
- Whiton RS, Witherspoon CL, Buckley TJ. Improved high-performance liquid chromatographic method for the determination of polycyclic aromatic hydrocarbon metabolites in human urine. *J Chromatogr B Biomed Appl* 1995;665(2):390-4.
- Wilson HK. Related biological monitoring values for occupational exposure: a United Kingdom perspective. *Int Arch Occup Environ Health* 1999;Jul;72(4):274-8.
- Woodruff RS. Confidence intervals for medians and other position measures. *Journal of the American Statistical Association* 1952; 47:635-47.
- World Health Organization. Environmental Health Criteria documents. [online]. Available from URL: http://www.who.int/pcs/ra_site/ehc.html, 9/30/02.
- Xu X, Duncan AM, Wangen KE, Kurzer MS. Soy consumption alters endogenous estrogen metabolism in postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2000;9(8):781-6.
- Zschiesche W, Schaller KH, Weltle D. Exposure to soluble barium compounds: an interventional study in arc welders. *Int Arch Occup Environ Health* 1992;64(1):13-23.

Appendix A

References for Biomonitoring Analytical Methods

Dioxins, Furans, PCBs, Organochlorine Pesticides

Turner W, DiPietro E, Cash TP, McClure PC, Patterson, DG Jr., Shirkhan H. An improved SPE extraction and automated sample cleanup method for serum PCDDs, PCDFs, and coplanar PCBs. *Organohalogen Compounds* 1994;19:31-5.

Turner W, DiPietro E, Lapeza C, Green V, Gill J, Patterson, DG Jr, et al. Universal automated cleanup system for the isotope-dilution high-resolution mass spectrometric analysis of PCDDs, PCDFs, coplanar PCBs, PCB congeners, and persistent pesticides from the same serum sample. *Organohalogen Compounds* 1997;31:26-31.

Metals

Chen HP, Paschal DC, Miller DT, Morrow JC. Determination of total and inorganic mercury in whole blood by on-line digestion with flow injection. *Atomic Spectroscopy* 1998;19:176-9.

Miller, DT, Paschal DC, Gunter EW, Stroud PE, D'Angelo J. Determination of lead in blood using electrothermal atomization atomic absorption spectrometry with a L'vov platform and matrix modifier. *Analyst* 1987;112:1701-4.

Paschal DC, Ting BG, Morrow JC, Pirkle JL, Jackson RJ, Sampson EJ, et al. Trace metals in urine of United States residents: reference range concentrations. *Environ Res* 1998;76:53-9.

Stoeppler M, Brandt K. Determination of cadmium in whole blood and urine by electrothermal atomic-absorption spectrophotometry. *Fresnus A Anal Chem* 1980;300:372-80.

Pesticide Metabolites

Beeson MD, Driskell WJ, Barr DB. Isotope dilution high-performance liquid chromatography/tandem mass spectrometry method for quantifying urinary metabolites of atrazine, malathion, and 2,4-dichlorophenoxyacetic acid. *Anal Chem* 1999;71(16):3526-30.

Bravo R, Driskell WJ, Whitehead RD Jr, Needham LL, Barr DB. Quantitation of dialkyl phosphate metabolites of organophosphate pesticides in human urine using GC-MS-MS with isotopic internal standards. *J Anal Toxicol* 2002;26:245-52.

Hill RH Jr, Shealy DB, Head SL, Williams CC, Bailey SL, Gregg M, et al. Determination of pesticide metabolites in human urine using isotope dilution technique and tandem mass spectrometry. *J Anal Toxicol* 1995;19(5):323-9.

PAHs

Smith CJ, Huang WL, Walcott CJ, Turner W, Grainger J, Patterson DG Jr. Quantification of monohydroxy-PAH metabolites in urine by solid-phase extraction with isotope dilution GC-HRMS. *Analytical and Bioanalytical Chemistry* 2002; 372:216-20.

Phthalate Metabolites

Blount BC, Milgram KE, Silva M, Malek N, Reidy J, Needham LL, et al. Quantitative detection of eight phthalate metabolites in human urine using HPLC-APCI-MS/MS. *Anal Chem* 2000;72:4127-34.

Phytoestrogens

Valentin-Blasini L, Blount BC, Rogers HS, Needham LL. HPLC-MS/MS method for the measurement of seven phytoestrogens in human serum and urine. *J Expo Anal Environ Epidemiol* 2000;10:799-807.

Tobacco Smoke (Cotinine)

Bernert JT, Turner WE, Pirkle JL, Sosnoff CS, Akins JR, Waldrep MK, et al. Development and validation of a sensitive measurement of serum cotinine in both smokers and nonsmokers by liquid chromatography/atmospheric pressure ionization tandem mass spectrometry. *Clin Chem* 1997;43:2281-91.

Appendix B

Confidence Interval Estimation for Percentiles

A common practice to calculate confidence intervals from survey data is to use large-sample normal approximations. 95% confidence intervals on point estimates of percentiles are often computed by adding and subtracting from the point estimate a quantity equal to twice its standard error. This normal approximation method may not be adequate, however, when estimating the proportion of subjects above or below a selected value (especially when the proportion is near 0.0 or 1.0 or when the effective sample size is small). In addition, confidence intervals on proportions deviating from 0.5 are not theoretically expected to be symmetric around the point estimate. Also, adding and subtracting a multiple of the standard error to an estimate near 0.0 or 1.0 can lead to impossible confidence limits (i.e., proportion estimates below 0.0 or above 1.0).

We used the method of Korn et al. (1998) to obtain 95% confidence intervals about percentile estimates. We describe the method below, using SAS Proc Univariate and SUDAAN.

Procedure to calculate confidence intervals about percentiles.

Step 1: Use SAS (SAS Institute Inc. 1999) Proc Univariate to obtain a point estimate of the percentile of a chemical's results for the demographic group of interest, for example, the 90th percentile of blood lead results for children aged 1-5 years. Use the Freq option to assign the correct sample weight for each chemical result.

Step 2: Use SUDAAN (SUDAAN Users Manual, 2001) Proc Descript with Jackknife design option to estimate the proportion of subjects with results below the percentile estimate obtained in Step 1 and to obtain the standard error associated with this proportion estimate.

Step 3: After obtaining an estimate of the proportion of subjects below the estimated percentile value (i.e., the proportion obtained in Step 2), compute the Clopper-Pearson 95% confidence interval using the degrees-of-freedom adjusted effective sample size (Korn and Graubard, 1998). (Note: If the degrees-of-freedom adjusted effective sample size is greater than the actual sample size, the actual sample size should be used.) This step will produce a lower and an upper limit for the estimated proportion obtained in Step 2.

Step 4: Use SAS Proc Univariate (again using the Freq option to assign weights) to determine the chemical values that correspond to the proportion obtained in Step 2 and the lower and upper limits on this proportion obtained in Step 3.

Example:

To estimate the 75th percentile, use SAS Proc Univariate with the Freq option to get a point estimate of the chemical value that corresponds to the 75th percentile. Then use SUDAAN to estimate the proportion of subjects with results below the 75th percentile (which should be very near 0.75). Next, obtain a confidence interval on this proportion by computing the Clopper-Pearson 95% confidence limits using the degrees-of-freedom adjusted effective sample size. Suppose these confidence limits are 0.67 and 0.81, then use SAS Proc Univariate with the Freq option to determine the chemical values corresponding to the 67th and 81st percentiles. These point estimates are the lower and upper confidence limits on the 75th percentile.